

Subgroups of R. Thompson's Group F that are Isomorphic to F .

Bronlyn Wassink

ABSTRACT:

This paper studies when a pair of elements in F are the images of the standard generators of F under a self monomorphism.

1 Introduction

Richard Thompson's Groups $F \subseteq T \subseteq V$ were defined by Richard Thompson in 1965. Thompson proved that both T and V are finitely presented infinite simple groups. The group F is also finitely presented, but it is not simple. A good introduction to these groups is in [9].

In this paper we find all subgroups of F isomorphic to F by finding each pair of elements of F that is the image, under a monomorphism of F into itself, of a standard pair of generators of F . This paper was motivated by the questions about the distortion of subgroups of F . It has been shown by José Burillo in [7] that for any natural number n , there is a subgroup of F that is isomorphic to $F^n \times \mathbb{Z}$ that can be quasi-isometrically embedded into F using the inclusion function. It is unknown if every subgroup of F that is isomorphic to F or $F \times \mathbb{Z}$ has this property. In order to get started on this question, we must first find every subgroup of F that is isomorphic to F .

There are several ways to define Thompson's Groups. The definition that this paper will focus on defines F as a subgroup of $\text{PL}_0(\mathbf{I})$, which is the group of all piecewise-linear, orientation preserving homeomorphisms of the unit interval, $\mathbf{I}$, that admit only finitely many points of non-differentiability. The group F is exactly the elements of $\text{PL}_0(\mathbf{I})$ whose points of non-differentiability all occur at dyadic rationals and have all slopes integral powers of 2. The dyadic rationals are defined to be the rational numbers of the form $\frac{m}{2^n}$, where $m, n \in \mathbb{Z}$ and $n > 0$.

In the first section of this paper, many standard results and terminology will be introduced.

The second section of this paper gives our main result. The presentation $\langle x_0, x_1 \mid [x_0 x_1^{-1}, x_1^{x_0}] = [x_0 x_1^{-1}, x_1^{x_0}] = 1 \rangle$ is called the "standard" finite presen-

tation for F . We find all pairs (f, g) of elements in $\text{PL}_0(\mathbf{I})$ for which there is an isomorphic embedding of F into $\text{PL}_0(\mathbf{I})$ that carries x_0 to f and x_1 to g .

The third section of this paper gives a partial converse to the main theorem in [3]. This section also provides a counterexample to the full converse of the main result in [3].

In addition to the application discussed in Section 3, the results in Section 2 of this paper are used in the complete classification of the isomorphism classes of the finite index subgroups of F . This classification appears in [2], which is a joint paper of the author and Collin Bleak. Independently from our work, Burillo, Cleary and Röver in [8] also characterized the finite index subgroups of F that are isomorphic to F as a result of their investigations into the commensurator of F . Their techniques are very different from those in [2]

1.1 Definitions and Notation

Richard Thompson's Group F can also be described by the following presentations.

$$F \cong \langle x_0, x_1, x_2, \dots \mid x_j^{x_i} = x_{j+1} \text{ for } i < j \rangle \quad (1)$$

$$F \cong \langle x_0, x_1 \mid [x_0 x_1^{-1}, x_1^{x_0}] = [x_0 x_1^{-1}, x_1^{x_0^2}] = 1 \rangle \quad (2)$$

where $a^b = b^{-1}ab$ and $[a, b] = aba^{-1}b^{-1}$.

Define the *standard finite presentation of F* to be the presentation (2) above.

We will make extensive use of the representation of F in the group $\text{PL}_0(\mathbf{I})$, whose definition is given in the introduction.

Composition and evaluation of functions in F will be in word order. That is, if $f, g \in F$ and $t \in [0, 1]$, then $tf = f(t)$, $fg = g \circ f$, and $f^g = g^{-1}fg = g \circ f \circ g^{-1}$.

The functions x_0 and x_1 are defined below. The standard representation of F as PL homeomorphisms of $\mathbf{I}$ carries x_0 and x_1 of (2) to the functions below with the same names.

$$ax_0 = \begin{cases} 2a & 0 \leq a \leq 1/4 \\ a + 1/4 & 1/4 \leq a \leq 1/2 \\ a/2 + 1/2 & 1/2 \leq a \leq 1 \end{cases} \quad ax_1 = \begin{cases} a & 0 \leq a \leq 1/2 \\ 2a - 1/2 & 1/2 \leq a \leq 5/8 \\ a + 1/8 & 5/8 \leq a \leq 3/4 \\ a/2 + 1/2 & 3/4 \leq a \leq 1 \end{cases}$$

One can check, using the conventions above, that the functions x_0 and x_1 satisfy the relations in (2). Presentations (1) and (2) are equivalent (see [9] Theorem 3.4). The fact that the functions x_0 and x_1 generate all of the claimed functions in F (as a group of homeomorphisms) is Corollary 2.6 in [9]. (Note that our functions x_0 and x_1 are the inverses of the homeomorphisms in [9].)

Given a homeomorphism $f : [0, 1] \rightarrow [0, 1]$, we will denote by $\text{Supp}(f)$ the *support* of f , defined as

$$\text{Supp}(f) = \{x \in [0, 1] \mid xf \neq f\}.$$

Given any $f \in \text{PL}_0(\mathbf{I})$, $\text{Supp}(f)$ will be a finite union of disjoint open intervals. Each of these open intervals will be called an *orbital* of f .

As $F \leq \text{PL}_0(\mathbf{I})$, many lemmas and theorems will be done using more general functions from $\text{PL}_0(\mathbf{I})$ instead of the more restrictive functions in F .

We will say that two functions f_0 and f_1 in $\text{PL}_0(\mathbf{I})$ generate a *standard isomorphic copy* of F if the subgroup of $\text{PL}_0(\mathbf{I})$ that is generated by f_0 and f_1 is isomorphic to F by an isomorphism sending x_0 to f_0 and x_1 to f_1 . For f_0 and f_1 to generate a standard isomorphic copy of F , it is necessary that $[f_0 f_1^{-1}, f_1^{f_0}] = [f_0 f_1^{-1}, f_1^{f_0^2}] = 1$.

1.2 Previous Results

The first remark below, though trivial, is useful in several proofs. It follows because the relations in the remark are relations in (2).

Remark 1.2.1 *If f_0 and f_1 are functions in $\text{PL}_0(\mathbf{I})$ that satisfy both $[f_0 f_1^{-1}, f_1^{f_0}] = 1$ and $[f_0 f_1^{-1}, f_1^{f_0^2}] = 1$, then there is a group homomorphism from F to $\langle f_0, f_1 \rangle$ sending x_0 to f_0 and x_1 to f_1 . In particular, f_0 and f_1 will satisfy all of the relations that x_0 and x_1 they will satisfy in F , such as $[f_1 f_2^{-1}, f_2^{f_1}] = 1$, where $f_2 = f_1^{f_0}$.*

Lemma 1.2.2 *If $f, g \in PL_0(\mathbf{I})$ where the support of f is disjoint from the support of g , then f and g commute.*

Lemma 1.2.3 *Let $g_1, g_2, \dots, g_n \in PL_0(\mathbf{I})$. Let H be the subgroup of $PL_0(\mathbf{I})$ that is generated by $g_1, g_2, \dots, g_n$ and define*

$$\text{Supp}(H) = \{x \in [0, 1] | xh \neq x \text{ for some } h \in H\}.$$

Then, $\text{Supp}(H) = \bigcup_{i=1}^n \text{Supp}(g_i)$.

Whenever H is a subgroup of $PL_0(\mathbf{I})$, the support of H will consist of disjoint open intervals. Call each of those intervals an *orbital* of H .

The following Lemma is a restatement of Remark 2.1, Lemma 2.2 and Lemma 2.5 in [1].

Lemma 1.2.4 1. *Any element $h \in PL_0(\mathbf{I})$ has only finitely many orbitals.*

2. *Let h be a function in $PL_0(\mathbf{I})$. If A is an orbital for h , then either $xh > x$ or $xh < x$ for all points x in A .*

3. *If $h \in PL_0(\mathbf{I})$ and $A = (a, b)$ is an orbital of h , then given any $\varepsilon > 0$ and x in A , there is an integer n so that $|xh^{-n} - a| < \varepsilon$ and $|xh^n - b| < \varepsilon$.*

4. *Let $g, h \in PL_0(\mathbf{I})$. (a, b) is an orbital of g if and only if (ah, bh) is an orbital of g^h .*

5. *If (a, b) is an orbital of f , $c \in (a, b)$, and $m \in \mathbb{Z}$, then $cf^m \in (a, b)$.*

Lemma 1.2.4 allows us to introduce new terminology. From Lemma 1.2.4.(1), since there are only finitely many orbitals, we can introduce an order on them. If A and B are orbitals of g , then we say that $A < B$ when $a < b$ for any $a \in A$ and $b \in B$. With this ordering, we can refer to the *first orbital*, the *second orbital*, and so on. Many times we will be interested in the first orbital and the last orbital.

Lemma 1.2.4.(2) allows us to introduce a sign on the orbitals. If an orbital A of g has the property that $ah > a$ for all a in A , then we call A an *up-bump* of g . Similarly, if an orbital B of g has the property that $bh < b$ for all b in B , then we call B a *down-bump* of g . Notice that Lemma 1.2.4.(2) guarantees that every orbital is either an up-bump or a down-bump.

We call the orbitals (a, b) of g and (ah, bh) of g^h *corresponding orbitals*.

The following lemma is a part of Lemma 2.5 in [1].

Lemma 1.2.5 *If $f, g \in PL_0(\mathbf{I})$ and (a, b) is an orbital of f , then the derivative from the right of f at a equals the derivative from the right of f^g at ag and the derivative from the left of f at b equals the derivative from the left of f^g at bg .*

In other words, f^g has the same *leading* and *trailing slopes* on its orbitals as f has on each of its corresponding orbitals.

Remark 1.2.6 *Given any functions f_0 and f_1 in $PL_0(\mathbf{I})$ and $k \geq 1$, we can define $f_{k+1} = f_1^{f_0^k}$. So then all of the f_k for $k \geq 1$ have the same leading slopes on each of their corresponding orbitals, and all f_k have the same trailing slopes on their corresponding orbitals.*

The following is Theorem 4.3 from [9].

Lemma 1.2.7 *F has no proper non-abelian quotients.*

In particular, if we can find an f and g where

$$[fg^{-1}, g^f] = [fg^{-1}, g^{f^2}] = 1$$

and f and g do not commute, then f and g generate a group that is isomorphic to F .

The following lemma is a combination of Theorems 4.1 and 4.5 in [9].

Lemma 1.2.8 *The group $F' = [F, F]$, the commutator subgroup of F , is simple. Furthermore, $\text{Supp}(F') = (0, 1)$ and F' consists of all of the functions $f \in F$ such that both $f'(0) = 1$ and $f'(1) = 1$, where $f'(0)$ is the derivative of f at zero from the right and $f'(1)$ is the derivative of f at one from the left.*

The following is a result by Brin and Squier in Theorem 4.18 in [6].

Theorem 1.2.9 *If A is an orbital of both f and g in $PL_0(\mathbf{I})$ and $[f, g]|_A = 1$, then there is a function h and integers m and n so that $h^m|_A = f|_A$ and $h^n|_A = g|_A$.*

2 Standard Isomorphic Copies of F .

In this section, we characterize those pairs (f, g) of elements in $\text{PL}_0(\mathbf{I})$ that generate a standard isomorphic copy of F .

This section will be organized as follows. Section 2.1 gives information about commuting functions. Then, assuming that a pair (f_0, f_1) of functions in $\text{PL}_0(\mathbf{I})$ generate a standard isomorphic copy of F , Sections 2.2 and 2.3 accumulate a series of increasingly restrictive observations extracted from the fact that f_0 and f_1 must satisfy the relations in (2). These give a set of conditions that are necessary for (f_0, f_1) to generate a standard isomorphic copy of F , but they will not quite be sufficient. They are sufficient however to imply that f_0 and f_1 satisfy the second relation in (2). Sections 2.4 and 2.5 then find the remaining conditions needed to characterize when (f_0, f_1) generate a standard isomorphic copy of F .

In the introduction we remark that for f_0 and f_1 in $\text{PL}_0(\mathbf{I})$ to generate a standard isomorphic copy of F it is necessary that $[f_0 f_1^{-1}, f_1^{f_0}] = [f_0 f_1^{-1}, f_1^{f_0^2}] = 1$ hold. By Lemma 1.2.7, we see that this is a sufficient condition when f_0 and f_1 do not commute.

If f_0 and f_1 in $\text{PL}_0(\mathbf{I})$ generate a standard isomorphic copy of F , we define $f_{k+1} = f_1^{f_0^k}$ for all $k \geq 0$.

2.1 Commuting Elements

Lemma 2.1.1 *Let $f, g \in \text{PL}_0(\mathbf{I})$ where $[f, g] = 1$. If A is an orbital of f , either A is disjoint from the support of g or A is also an orbital of g .*

Proof: Suppose $fg = gf$. Let $A = (a, c)$ be an orbital of f that is not disjoint from the support of g . Let B be an orbital of g where $A \neq B$ and $A \cap B \neq \emptyset$. By these assumptions, either f has a fixed point in B or g has a fixed point in A . Without loss of generality, we may assume that g has a fixed point in A and that A is an up-bump of f .

Suppose that $y \in (a, c)$ is a fixed point of g . For all $k \in \mathbb{Z}$ we have $y f^k g = y g f^k = y f^k$. So $y f^k$ is in the interval (a, c) and is not in an orbital of g . Since g can only have finitely many orbitals and $\lim_{k \rightarrow -\infty} b f^k = a$, then there must be some interval (a, t) that is disjoint from the support of g . Because there are points in (a, c) that are in the support of g , then there will be an orbital (b, d) of g where $b \in (a, c)$ and (a, b) is disjoint from the support of

g . Let $x \in (b, fb) \cap (b, d)$. Let $z = xf^{-1}$. So z is a fixed point of g and zf is not a fixed point of g . Then $(zg)f = zf \neq (zf)g$, a contradiction. $\square$

Lemma 2.1.2 *Let $f_0, f_1 \in PL_0(\mathbf{I})$ such that $[f_1^{f_0}, f_0f_1^{-1}] = 1$ and $[f_0f_1^{-1}, f_1^{f_0^2}] = 1$. If A is an orbital of both f_0 and f_1 , then f_0 and f_1 commute on A .*

Proof: Suppose $A = (a, c)$ is an orbital of both f_0 and f_1 . Since $f_{k+1} = f_1^{f_0^k}$, $(af_0^k, cf_0^k) = (a, c) = A$ is also an orbital for all f_{k+1} , $k \geq 1$. Note that $[f_1f_2^{-1}, f_1^{-1}f_2f_1] = 1$. By Remark 1.2.6, the leading slopes on A of f_1 and f_2 are equal and the trailing slopes on A of f_1 and f_2 are equal. So there is an $a' \neq a$ with $a \in A$ and a $c' \neq c$ with $c \in A$ such that $f_1|_{[a, a']} = f_2|_{[a, a']}$ and $f_1|_{[c', c]} = f_2|_{[c', c]}$. Then $f_1f_2^{-1}$ is the identity on $[a, a'] \cup [c', c]$. So then any orbital of $f_1f_2^{-1}$ is either disjoint from $[a, c]$ or is contained in $[a', c']$. Since f_3 and $f_1f_2^{-1}$ commute, by Lemma 2.1.1 the orbitals of these functions must either be disjoint or equal. If $P \subseteq [a', c']$ is an orbital $f_1f_2^{-1}$, then P and (a, c) can neither be disjoint nor equal. So then $P = \emptyset \Rightarrow (f_1f_2^{-1})|_A = 1 \Rightarrow f_1|_A = f_2|_A$. Thus $(f_0^{-1}f_1f_0)|_A = f_2|_A = f_1|_A \Rightarrow (f_1f_0)|_A = (f_0f_1)|_A$. $\square$

2.2 Orbital Arrangements

Lemma 2.2.1 *Let $f_0, f_1 \in PL_0(\mathbf{I})$ such that $[f_1^{f_0}, f_0f_1^{-1}] = 1$ and $[f_0f_1^{-1}, f_1^{f_0^2}] = 1$. Let $A_1, A_2, \dots, A_N$ be the orbitals of f_0 . Let $B_1, B_2, \dots, B_M$ be the orbitals of f_1 . Then for any A_j and B_i where $A_j \cap B_i \neq \emptyset$, either $A_j \subseteq B_i$ or $B_i \subseteq A_j$.*

Proof: Suppose there is an A_j and a B_i such that $A_j \cap B_i \neq \emptyset$ but $A_j \not\subseteq B_i$ and $B_i \not\subseteq A_j$. Let $A_j = (a, c)$ and $B_i = (b, d)$. Either $a < b < c < d$ or $b < a < d < c$. Assume that $a < b < c < d$. One of the orbitals of $f_1^{f_0}$ is (bf_0, df_0) . Since $b < c < d$, then $bf_0 < cf_0 < df_0$. Since c is a fixed point of f_0 , the previous inequality simplifies to $bf_0 < c < df_0$. So c is in an orbital of $f_1^{f_0}$.

Since c is not a fixed point of f_1 and is a fixed point of f_0 , $cf_0f_1^{-1} = cf_1^{-1} \neq c$. So c is not a fixed point of $f_0f_1^{-1}$. So c is in an orbital of $f_0f_1^{-1}$.

Since the orbital (bf_0, df_0) of $f_1^{f_0}$ is not disjoint from the set of all orbitals of $f_0f_1^{-1}$ (c is in their intersection), then, by Lemma 2.1.1, an orbital for $f_0f_1^{-1}$ must also equal (bf_0, df_0) .

The point c is also in the interval (bf_0^2, df_0^2) , which is an orbital for $f_1^{f_0^2}$. Since the orbitals of $f_0f_1^{-1}$ and $f_1^{f_0^2}$ are not disjoint (again, both contain the point c), then, by Lemma 2.1.1, an orbital for $f_0f_1^{-1}$ must equal (bf_0^2, df_0^2) .

Since b is not a fixed point for f_0 , then either $bf_0 < bf_0^2 < c$ or $bf_0^2 < bf_0 < c$. So (bf_0^2, df_0^2) and (bf_0, df_0) are non-equal, non-disjoint intervals. Thus they can not both be orbitals for the same function, $f_0f_1^{-1}$, which is a contradiction.

Similarly, if $b < a < d < c$, then an almost identical argument will show that a is in the orbital (bf_0, df_0) of $f_1^{f_0}$, the orbital (bf_0^2, df_0^2) of $f_1^{f_0^2}$, and also in an orbital of $f_0f_1^{-1}$. Thus by Lemma 2.1.1, these three orbitals must all be equal, which contradicts the fact that $(bf_0, df_0) \neq (bf_0^2, df_0^2)$. $\square$

Lemma 2.2.2 *Let $f_0, f_1 \in PL_0(\mathbf{I})$ such that $[f_1^{f_0}, f_0f_1^{-1}] = 1$ and $[f_0f_1^{-1}, f_1^{f_0^2}] = 1$. If A is an orbital of f_0 then either there exists an orbital B of f_1 where $B \subseteq A$, or A is disjoint from the support of f_1 .*

Proof: By Lemma 2.2.1, if $A \cap B \neq \emptyset$, then either $A \subseteq B$ or $B \subseteq A$. Suppose $A \not\subseteq B$. Let $B = (b, d)$ and let $A = (a, c)$. Suppose b is not a fixed point of f_0 . Then it must be in some orbital (r, s) of f_0 , where $r < b < s < a$. This contradicts Lemma 2.2.1. It is similarly shown that d is a fixed point of f_0 . So then $(bf_0, df_0) = (b, d) = B$ is an orbital of $f_0^{-1}f_1f_0 = f_2$. By Remark 1.2.1, $[f_2^{f_1}, f_1f_2^{-1}] = 1$ and $[f_1f_2^{-1}, f_2^{f_1^2}] = 1$. Then by Lemma 2.1.2, since B is an orbital for both f_1 and f_2 , we have that f_1 and f_2 must commute in B .

So $f_1f_2|_B = f_2f_1|_B$. Then $f_2|_B = f_1^{-1}f_2f_1|_B = f_3|_B = f_0^{-1}f_2f_0|_B$. So then $f_2f_0|_B = f_0f_2|_B$. So, by Lemma 2.1.1, either the orbitals A of f_0 and B of f_2 must either be disjoint or equal. But we assumed that $A \not\subseteq B$, which gives a contradiction. $\square$

2.3 Necessary Conditions

Lemma 2.3.1 *Let $f_0, f_1 \in PL_0(\mathbf{I})$ such that $[f_1^{f_0}, f_0f_1^{-1}] = 1$ and $[f_0f_1^{-1}, f_1^{f_0^2}] = 1$. Let $A = (a, c)$ be an orbital of f_0 that is an up-bump. Suppose that (a, c) is not an orbital of f_1 . Let $(b_1, d_1), (b_2, d_2), \dots, (b_n, d_n)$ be all of the orbitals of f_1 , in increasing order, where $(b_i, d_i) \cap (a, c) \neq \emptyset$. Then all of the following are true.*

- i. $a < b_1$.
- ii. There is a point $p < c$ such that $f_0|_{[p, c]} = f_1|_{[p, c]}$.
- iii. $d_n = c$.
- iv. If p is the minimal point where $f_0|_{[p, c]} = f_1|_{[p, c]}$, then $b_nf_0 \geq p$.
- v. $b_1f_0 > b_n$.

Proof: Recall from Lemma 2.2.2 that if B is an orbital of f_1 and (a, c) is an orbital of f_0 , then either $B \subseteq (a, c)$ or $B \cap (a, c) = \emptyset$. So then $a \leq b_1$ and $d_n \leq c$.

i. Suppose that $a = b_1$. Since (a, c) , is not an orbital of f_1 , we have $d_1 \neq c$. Since f_0 is an up-bump, $(a, d_1) \subset (a, d_1 f_0) \subset (a, d_1 f_0^2)$, where $(a, d_1 f_0)$ is an orbital for f_2 and $(a, d_1 f_0^2)$ is an orbital for f_3 . Recall that $f_2 = f_1^{f_0}$, $f_3 = f_2^{f_0}$, and $[f_2, f_0 f_1^{-1}] = [f_3, f_0 f_1^{-1}] = 1$. Since $d_1 f_1 f_0^{-1} = d_1 f_0^{-1} < d_1$, we see that d_1 is in some orbital of $f_0 f_1^{-1}$. So by Lemma 2.1.1, both $(a, d_1 f_0)$ and $(a, d_1 f_0^2)$ must be orbitals for $f_0 f_1^{-1}$. These are non-disjoint, non-equal intervals, thus they can not both be orbitals of $f_0 f_1^{-1}$, a contradiction.

ii. Suppose not. Then there is a number $t \in (a, c)$ so that (t, c) is an orbital of $f_0 f_1^{-1}$. Since c is an endpoint of an orbital of f_0 and $a < b_1$ (from *i*), it is possible to find an $k \in \mathbb{N}$ so that $t < b_1 f_0^k < d_1 f_0^k \leq c$. $(b_1 f_0^k, d_1 f_0^k)$ is an orbital for f_{k+1} , which commutes with $f_0 f_1^{-1}$. So then by Lemma 2.1.1, either the orbitals (t, c) and $(b_1 f_0^k, d_1 f_0^k)$ must be disjoint or equal, a contradiction.

iii. This follows immediately from *ii*.

iv. From *iii*, $(b_n f_0, c)$ is an orbital for f_2 . Since $f_0|_{[p, c]} = f_1|_{[p, c]}$, we know $[p, c]$ is disjoint from the orbitals of $f_0 f_1^{-1}$. Also, since p is minimal, there is a $w \in [a, p)$ so that (w, p) is an orbital of $f_0 f_1^{-1}$. By Lemma 2.1.1, it must be the case that $(b_n f_0, c)$ is disjoint from the orbitals of $f_0 f_1^{-1}$, giving $b_n f_0 \geq p$.

v. Suppose that $b_1 f_0 \leq b_n$. Recall that the first orbital of f_2 inside (a, c) is $(b_1 f_0, d_1 f_0)$. Since $f_1 f_k f_1^{-1} = f_{k+1}$ for all $k \geq 2$, the first orbital of f_{k+2} contained in (a, c) is $(b_1 f_0 f_1^k, d_1 f_0 f_1^k)$. Since $f_0 f_k f_0^{-1} = f_{k+1}$ for all $k \geq 2$, the first orbital of f_{k+2} contained in (a, c) is $(b_1 f_0 f_0^k, d_1 f_0 f_0^k)$. From the two previous statements, we see that for all $k \geq 2$, we have $b_1 f_0 f_0^k = b_1 f_0 f_1^k$. Since $b_1 \in (a, c)$, an orbital of f_0 , then there is a $k \in \mathbb{N}$ where $b_1 f_0 f_0^k > b_n$. Since $b_1 f_0 \leq b_n$, and, by *iii*, (b_n, c) is an orbital of f_1 , then for all $k \in \mathbb{N}$, we have $b_1 f_0 f_1^k \leq b_n$, which the contradiction that for appropriately large n , $b_n \geq b_1 f_0 f_1^k = b_1 f_0 f_0^k > b_n$. $\square$

Similarly, we get the analogous result for down bumps below.

Lemma 2.3.2 *Let $f_0, f_1 \in PL_0(\mathbf{I})$ such that $[f_1^{f_0}, f_0 f_1^{-1}] = 1$ and $[f_0 f_1^{-1}, f_1^{f_0^2}] = 1$. Let $A = (a, c)$ be an orbital of f_0 that is a down-bump. Assume that (a, c) is not an orbital of f_1 . Let $(b_1, d_1), (b_2, d_2), \dots, (b_n, d_n)$ be all of the orbitals of f_1 , in increasing order, where $(b_i, d_i) \cap (a, c) \neq \emptyset$. Then all of the following are true. *i.* $d_n < c$. *ii.* There is a point $\rho > a$ such that $f_0|_{[a, \rho]} = f_1|_{[a, \rho]}$. *iii.* $b_1 = a$. *iv.* If ρ is the maximal point where $f_0|_{[a, \rho]} = f_1|_{[a, \rho]}$, then $d_1 f_0 \leq \rho$.*

v. $d_n f_0 < d_1$.

Lemma 2.3.3 *Let f_0, f_1 be elements in $PL_0(\mathbf{I})$. Let (a, c) be an orbital of f_0 . Suppose that (a, c) is not an orbital of f_1 . Assume that whenever B is an orbital of f_1 , either $B \subseteq (a, c)$ or $B \cap (a, c) = \emptyset$. Let $(b_1, d_1), (b_2, d_2), \dots, (b_n, d_n)$ be all of the orbitals of f_1 , in increasing order, that are properly contained in (a, c) . If f_0 and f_1 either satisfy $i - v$ in Lemma 2.3.1 or $i - v$ in Lemma 2.3.2, then $[f_0 f_1^{-1}, f_1^{f_0^2}]|_{[a, c]} = 1$.*

Proof: Suppose that f_0 is an up-bump. So then f_0 and f_1 satisfy $i - v$ in Lemma 2.3.1. The first orbital of $f_1^{f_0^2}$ is $(b_1 f_0^2, d_1 f_0^2)$. By v, $b_1 f_0 > b_n$ so then $b_1 f_0 f_0 > b_n f_0 \geq p$ (by iv). So all of the orbitals of $f_1^{f_0^2}$ that are contained in (a, c) are in (p, c) . By ii, $f_0|_{[p, c]} = f_1|_{[p, c]}$, so then $f_0 f_1^{-1}|_{[p, c]} = 1$. Then the orbitals of $f_1^{f_0^2}|_{[a, c]}$ and $f_0 f_1^{-1}|_{[a, c]}$ are disjoint, thus the functions commute.

A similar argument shows that $[f_0 f_1^{-1}, f_1^{f_0^2}]|_{[a, c]} = 1$ when (a, c) is a down-bump of f_0 . $\square$

The lemma below characterizes the “nice” standard isomorphic copies of F , in that there are only 6 properties to check, all of which are completely determined by the function values of some key points of the domain. It is not a necessary condition that $b_1 f_0 \geq p$ (for up-bumps) or $d_1 f_0 \leq \rho$ (for down-bumps) in order for the functions f_0 and f_1 to satisfy the standard F relations.

Lemma 2.3.4 *Let f_0, f_1 be non-commuting functions in $PL_0(\mathbf{I})$ where for every orbital A of f_0 and B of f_1 , either $B \subseteq A$ or $A \cap B = \emptyset$. Assume that for every orbital (a, c) that is an up-bump of f_0 , if $(b_1, d_1), (b_2, d_2), \dots, (b_n, d_n)$ are all of the orbitals of f_1 , in increasing order, that are properly contained in (a, c) , then $i - v$ in Lemma 2.3.1 are satisfied as well as the additional condition $b_1 f_0 \geq p$. Also assume that for every orbital (a, c) that is a down-bump of f_0 , if $(b_1, d_1), (b_2, d_2), \dots, (b_n, d_n)$ are all of the orbitals of f_1 , in increasing order, that are properly contained in (a, c) , then $i - v$ in Lemma 2.3.2 are satisfied as well as the additional condition $d_n f_0 \leq \rho$. Then f_0 and f_1 generate a standard isomorphic copy of F .*

Proof: By Lemma 1.2.7, since f_0 and f_1 do not commute, it only needs to be shown that $[f_0 f_1^{-1}, f_1^{f_0}] = 1$ and $[f_0 f_1^{-1}, f_1^{f_0^2}] = 1$.

Since for every orbital A of f_0 and B of f_1 , either $B \subseteq A$ or $A \cap B = \emptyset$, then $[f_0 f_1^{-1}, f_1^{f_0}]|_{[0,1] - \text{Supp}(f_0)} = [f_1^{-1}, f_1]|_{[0,1] - \text{Supp}(f_0)} = 1$ and $[f_0 f_1^{-1}, f_1^{f_0^2}]|_{[0,1] - \text{Supp}(f_0)} = [f_1^{-1}, f_1]|_{[0,1] - \text{Supp}(f_0)} = 1$. Thus the relations in (2) only need to be checked in the orbitals of f_0 .

Let (a, c) be an orbital of f_0 . If (a, c) is an up-bump, then $p \leq b_1 f_0 < b_1 f_0^2 < c$. Recall that when f_0 is an up-bump in (a, c) , the part of $\text{Supp}(f_1^{f_0})$ contained in the interval (a, c) is equal to $(b_1 f_0, c)$ and the part of $\text{Supp}(f_1^{f_0^2})$ that is contained in the interval (a, c) is equal to $(b_1 f_0^2, c)$. This gives $\text{Supp}(f_1^{f_0})|_{(a,c)} \subseteq \text{Supp}(f_1^{f_0^2})|_{(a,c)} \subseteq (p, c)$ and $\text{Supp}(f_0 f_1^{-1}) \cap (p, c) = \emptyset$. So we have $[f_1^{f_0}, f_0 f_1^{-1}]|_{[a,c]} = 1$ and $[f_1^{f_0^2}, f_0 f_1^{-1}]|_{[a,c]} = 1$.

Similarly, when (a, c) is a down-bump of f_0 , then $a < d_n f_0^2 < d_n f_0 \leq \rho$. So then we have $\text{Supp}(f_1^{f_0^2})|_{(a,c)} \subseteq \text{Supp}(f_1^{f_0})|_{(a,c)} \subseteq (a, \rho)$ and $\text{Supp}(f_0 f_1^{-1}) \cap (a, \rho) = \emptyset$. So again we have $[f_1^{f_0^2}, f_0 f_1^{-1}]|_{[a,c]} = 1$ and $[f_1^{f_0}, f_0 f_1^{-1}]|_{[a,c]} = 1$. $\square$

2.4 Necessary and Sufficient Conditions

Previously in this paper, we have shown that there are several conditions that f_0 and f_1 must satisfy in order to generate a standard isomorphic copy of F . It has also been shown in Lemma 2.3.3 that these conditions are sufficient to guarantee that the relation $[f_0 f_1^{-1}, f_1^{f_0^2}] = 1$ holds. The lemmas in this section will focus on the relation $[f_0 f_1^{-1}, f_1^{f_0}] = 1$.

Lemma 2.4.1 *Assume f_0 and f_1 generate a standard isomorphic copy of F in $PL_0(\mathbf{I})$.*

Suppose that (a, c) is an up-bump of f_0 , (b_n, c) is an orbital of f_1 , and p is the minimal number so that $f_0|_{[p,c]} = f_1|_{[p,c]}$. If there is a number $q \in (a, p)$ so that $q f_0 = q f_1$, then $q \in (b_n, p)$.

Alternately, suppose that (a, c) is a down-bump of f_0 , (a, d_1) is an orbital of f_1 , and ρ is the minimal number so that $f_0|_{[a,\rho]} = f_1|_{[a,\rho]}$. If there is a number $q \in (\rho, c)$ so that $q f_0 = q f_1$, then $q \in (\rho, d_1)$.

Proof: In the case that (a, c) is an up-bump of f_0 , assume that $q \notin (b_n, p)$. Since q is in an orbital of f_0 , $q f_1 = q f_0 \neq q$. So then q must also be in some orbital, (b_k, d_k) , of f_1 . So then $q f_0 = q f_1 \in (b_k, d_k)$, where $d_k \leq b_n$. In particular, $q f_0 = q f_1 < d_k \leq b_n$. Combining the previous inequality with $b_k f_0 < q f_0$ and Lemma 2.3.1 part v , we get $q f_0 = q f_1 < d_k \leq b_n < b_1 f_0 \leq b_k f_0 < q f_0 = q f_1$, a contradiction.

The down-bump case is similar. $\square$

Lemma 2.4.2 *Assume f_0 and f_1 generate a standard isomorphic copy of F in $PL_0(\mathbf{I})$. Suppose that (a, c) is an orbital of f_0 that contains the orbitals $(b_1, d_1), \dots (b_n, d_n)$ of f_1 .*

Suppose that (a, c) is an up-bump of f_0 . If r is the minimal number in the interval (a, c) so that $rf_0 = rf_1$, then $b_1f_0 \geq r$.

Alternately, suppose that (a, c) is a down-bump of f_0 . If r is the maximal number in the interval (a, c) so that $rf_0 = rf_1$, then $d_nf_0 \leq r$.

Proof: In the case that (a, c) is an up-bump of f_0 , assume that r is the minimal number where $rf_0 = rf_1$. From Lemma 2.4.1, we have $r > b_n$. Since r is minimal, then (a, r) must be an orbital of $f_0f_1^{-1}$.

The interval (b_1f_0, d_1f_0) is the first orbital of $f_1^{f_0}$ contained in (a, c) . Since the orbitals of $f_0f_1^{-1}$ and $f_1^{f_0}$ must commute, then their orbitals must be disjoint or equal (Lemma 2.1.1).

As (a, r) can not equal the interval (b_1f_0, d_1f_0) , it follows that they must be disjoint, and $b_1f_0 \geq r$.

The case for down-bumps is similar. $\square$

Define a *nice orbital* of f_0 to be an orbital of f_0 where f_0 and f_1 satisfy the conditions of Lemma 2.3.4. For convenience, we may refer to *nice up-bumps* and *nice down-bumps* of f_0 .

Lemma 2.3.4 gives that any two functions f_0 and f_1 where every orbital of f_0 is either a nice orbital (or is an interval where f_0 and f_1 commute) will generate a standard isomorphic copy of F . We will call such a pair a *nice generating pair*.

We will reuse the notation in Lemmas 2.3.1, 2.4.1, and 2.4.2 to categorize the orbitals of f_1 into three disjoint sets.

1. The *Outside Orbitals*, which are orbitals contained in the interval (rf_0^{-1}, pf_0^{-1}) when (a, c) is an up-bump, and orbitals contained in the interval $(\rho f_0, rf_0)$ when (a, c) is a down-bump.
2. The *Inside Orbitals*, which are orbitals contained in the interval (pf_0^{-1}, b_n) when (a, c) is an up-bump, and orbitals contained in the interval $(d_1, \rho f_0)$ when (a, c) is a down-bump.
3. The *Main Orbital*, which is the orbital (b_n, c) when (a, c) is an up-bump and (a, d_1) when (a, c) is a down-bump.

Recall that Lemma 2.4.2 guarantees that for an up-bump, (b_1, pf_0^{-1}) is contained in the interval (rf_0^{-1}, pf_0^{-1}) . Also, for a down-bump, the same lemma gives that $(\rho f_0, d_n)$ is contained in $(\rho f_0, rf_0)$.

If f_1 does not have any Outside Orbitals in (a, c) , then (a, c) would be a nice orbital of f_0 , and all of the relations of F would be satisfied on this interval.

Theorem 2.4.3 *Let f_0 and f_1 be non-commuting functions in $PL_0(\mathbf{I})$ that satisfy $i - v$ in Lemmas 2.3.1 (for up-bumps) and 2.3.2 (for down bumps). Suppose also that every orbital of f_1 that is contained in an orbital of f_0 is either a Main Orbital, an Inside Orbital, or an Outside Orbital. The functions f_0 and f_1 generate a standard isomorphic copy of F if and only if every orbital of $f_1^{f_0}$ that corresponds to an Outside Orbital of f_1 commutes with $f_0 f_1^{-1}$.*

Proof:

($\Rightarrow$) We have shown in several previous lemmas that if f_0 and f_1 generate a standard isomorphic copy of F , then all of the listed properties must be true.

($\Leftarrow$) Since f_0 and f_1 are non-commuting, if they satisfy the relations of F then they generate a standard isomorphic copy of F .

As all orbitals of f_1 satisfy $i - v$ of 2.3.1 or 2.3.2, we have shown that these properties alone are enough to guarantee the second relation, $[f_0 f_1^{-1}, f_1^{f_0^2}] = 1$, holds.

Every orbital of $f_1^{f_0}$ that corresponds to a Main Orbital or an Inside Orbital of f_1 is disjoint from the support of $f_0 f_1^{-1}$, thus they commute. In addition, since every orbital of f_1 that does not commute with f_0 is an Outside Orbital and every orbital of $f_1^{f_0}$ that corresponds to an Outside Orbital of f_1 also commutes with $f_1 f_1^{-1}$, then we get the first relation $[f_0 f_1^{-1}, f_1^{f_0}] = 1$. $\square$

2.5 Building a Standard Generating Pair

We will end this section with a description of exactly how to construct any pair of functions, f_0 and f_1 , that generate a standard isomorphic copy of F .

To make an arbitrary pair of functions that generate a standard isomorphic copy of F , we will begin with two functions f_0 and f_1 where every orbital of f_0 is either a nice orbital or is an interval where f_0 and f_1 commute. We will then slightly modify the functions in a nice orbital of f_0 to construct

two new functions that contain Outside Orbitals of f_1 and also generate a standard isomorphic copy of F .

First, we will explicitly state how to build any nice generating pair f_0 and f_1 , where f_0 has exactly one nice up-bump. The process is almost identical for constructing pairs with exactly one nice down-bump, as well as constructing pairs with more than one nice orbital.

Remark 2.5.1 *Let f_0 be any function in $PL_0(\mathbf{I})$ that is not the identity function. In any up-bump (a, c) of f_0 , pick any point $p \in (a, c)$. Let f_1 be any function in $PL_0(\mathbf{I})$ where $f_0 = f_1$ on the interval $[p, c]$, where f_1 is the identity in the interval (a, pf_0^{-1}) , and where f_1 and f_0 commute outside of (a, c) . The functions f_0 and f_1 generate a standard isomorphic copy of F .*

Theorem 2.5.2 *Assume that (a, c) is a nice up-bump of f_0 . Let p be as in Lemma 2.3.4. Let $s \in (p, c)$. If h be any function in $PL_0(\mathbf{I})$ where $\text{Supp}(h) \subseteq (pf_0^{-1}, sf_0^{-1})$ and t, k are any integers, then the pair f_0 and f_1 are a standard generating pair of F if and only if f_0 and $g_1 = h^t f_0^{-1} h^k f_0 f_1$ are a standard generating pair of F .*

Proof: As the functions f_1 , and g_1 are identical outside of the interval (a, c) , then the pair f_0 and f_1 satisfy the relations of F outside the interval (a, c) if and only if the relations are also satisfied outside of (a, c) by the pair f_0 and g_1 . As f_0 and f_1 satisfy the conditions of Lemma 2.3.4, then if f_0 and g_1 are a standard generating pair, then so are f_0 and f_1 .

Suppose that f_0 and f_1 are a standard generating pair for F . By construction, f_0 and g_1 satisfy the conditions in Lemma 2.3.3, thus only the first relation must be checked for these functions. Using the fact that $f_0 f_1^{-1}$ is the identity on the interval (p, c) , it is a straightforward to check that whenever $y \in (p, s)$, we have $y f_0 g_1^{-1} g_1^{f_0} = y g_1^{f_0} f_0 g_1^{-1} = y f_0^{-1} h^{t-k} f_0$. Combine this computation with the facts that $g_1^{f_0}$ is the identity in the interval (a, p) and $f_0 g_1^{-1}$ is the identity in the interval (s, c) to get that the relation $[g_1^{f_0}, f_0 g_1^{-1}] = 1$ holds. $\square$

Theorem 2.5.3 *If f_0 and g_1 are a standard generating pair of F , then f_0 and g_1 can be inductively constructed from a nice generating pair f_0, f_1 in a finite number of steps as described in Theorem 2.5.2 (or the theorem's analogous result for down-bumps).*

Proof: Assuming that f_0 and g_1 are not a nice generating pair of F , there must be an orbital (a, c) of f_0 that properly contains an Outside Orbital of g_1 . Without loss of generality, assume (a, c) is an up-bump. Let p be the minimal point where $f_0|_{[p, c]} = g_1|_{[p, c]}$. Let α be the minimal point in (a, c) where $\alpha f_0 = \alpha g_1$. The first orbital $(b_1 f_0, d_1 f_0)$ of $g_1^{f_0}$ in (a, c) must either be an orbital of $f_0 g_1^{-1}$ or disjoint from the support of $f_0 g_1^{-1}$. In either case, $(b_1 f_0) f_0 = (b_1 f_0) g_1$, so then $\alpha \leq b_1 f_0 < p$. From Lemma 2.4.1, $b_n < \alpha$.

Define f_1 to be the function in $\text{PL}_0(\mathbf{I})$ that is equal to g_1 outside of the interval (a, c) , is the identity function in the interval $(a, p f_0^{-1})$, is equal to g_1 in the interval $(p f_0^{-1}, \alpha)$, and is equal to f_0 in the interval (α, c) . As all the conditions of Lemma 2.3.4 are satisfied for f_0 and f_1 , the pair f_0 and f_1 are a standard generating pair for F , and has one less non-nice orbital than f_0 and g_1 .

Let $(b_1, d_1), \dots, (b_m, d_m)$ be the orbitals of g_1 that are not orbitals of f_1 . Let $(q_1, w_1), \dots, (q_z, w_z)$ be the orbitals of $f_0 g_1^{-1}$ that are contained in (α, c) . As there are no non-equal, overlapping intervals in the set $S = \{(b_i f_0, d_i f_0), (q_j, w_j)\}$, we will always look at the first orbital in the set. This will give three cases for each step in the inductive process. Let $g_{(1,0)} = f_1$. The following three cases will construct $g_{(1,1)}$.

For the first case, assume $(b_1 f_0, d_1 f_0)$ is the unique first interval in the set S . Then let h_1 be the function in $\text{PL}_0(\mathbf{I})$ where $\text{Supp}(h_1) \subseteq (b_1, d_1)$ and $h_1|_{(b_1, d_1)} = g_1|_{(b_1, d_1)}$. Let $g_{(1,1)} = h_1^1 f_0^{-1} h_1^0 f_0 g_{(1,0)} = h_1 f_1$, which equals g_1 on (b_1, d_1) and $(p f_0^{-1}, d_1 f_0)$. Now remove $(b_1 f_0, d_1 f_0)$ from the set S and relabel $(b_i f_0, d_i f_0)$ in S to $(b_{i-1} f_0, d_{i-1} f_0)$.

For the second case, assume that (q_1, w_1) is the unique first interval in the set S . Then let ψ be the function in $\text{PL}_0(\mathbf{I})$ where $\text{Supp}(\psi) \subseteq (q_1, w_1)$ and $\psi|_{(q_1, w_1)} = f_0 g_1^{-1}|_{(q_1, w_1)}$. Let $h_1 = f_0 \psi^{-1} f_0^{-1}$. Then if we let $g_{(1,1)} = h_1^0 f_0^{-1} h_1^1 f_0 g_{(1,0)}$, we have $g_{(1,1)}$ equals g_1 on $(q_1 f_0^{-1}, w_1 f_0^{-1})$ and $(p f_0^{-1}, w_1)$. Now remove (q_1, w_1) from the set S and relabel (q_j, w_j) in S to (q_{j-1}, w_{j-1}) .

For the third case, assume $(b_1 f_0, d_1 f_0) = (q_1, w_1)$. Then by Theorem 1.2.9, $g_1^{f_0}|_{(b_1 f_0, d_1 f_0)}$ and $f_0 g_1^{-1}|_{(b_1 f_0, d_1 f_0)}$ are powers of a common root function. Let ϕ be the function in $\text{PL}_0(\mathbf{I})$ whose support is contained in $(b_1 f_0, d_1 f_0)$ and t and $-k$ be the integers so that on the interval $(b_1 f_0, d_1 f_0)$, we have $\phi^t = g_1 f_0$ and $\phi^{-k} = f_0 g_1^{-1}$. In this case, let $h_1 = \phi^{f_0^{-1}}$. Then the function $g_{(1,1)} = h_1^t f_0^{-1} h_1^k f_0 g_{(1,0)}$ is equal to f_1 on the intervals (b_1, d_1) and $(p f_0^{-1}, d_1 f_0)$. Now remove both $(b_1 f_0, d_1 f_0)$ and (q_1, w_1) from S and relabel $(b_i f_0, d_i f_0)$ to $(b_{i-1} f_0, d_{i-1} f_0)$ and (q_j, w_j) to (q_{j-1}, w_{j-1}) .

In each of the three cases, S becomes smaller by one or two intervals. For the β^{th} step in the inductive process, find the function h_β and integers r, k as in steps 1 – 3 above, and let $g_{(1,\beta)} = h_1^t f_0^{-1} h_1^k f_0 g_{(1,\beta-1)}$. After a finite amount of steps, S will become empty and the last function created, $g_{(1,\sigma)}$, will be equal to g_1 . $\square$

3 Partial Converse of the Ubiquity Result

The following theorem is the Ubiquity result, which is the main theorem in [3].

Theorem 3.1 *If H is a subgroup of $PL_0(\mathbf{I})$ and there exists an orbital $W = (a, b)$ of H and a function f in H so that either $f|_W$ moves points near a but not b or $f|_W$ moves points near b but not a , then there is a subgroup G of H so that G is isomorphic to F .*

A partial converse to the Ubiquity result has previously been proven by Guba in an unpublished paper, whose proof used much more machinery than what was used in this thesis. A proof with significantly less machinery is given below.

Theorem 3.2 *If $F \cong H \leq PL_0(\mathbf{I})$, then there exists an orbital $W = (a, c)$ of H and a function h in H where either $h|_W$ moves points near a but not c or $h|_W$ moves points near c but not a .*

Proof: Since $F \cong G$, there are functions f_0 and f_1 in H that generate a standard isomorphic copy of F . Then by Theorem ??, there exists an orbital (a, c) of f_0 that properly contains an orbital of f_1 . From Theorem ?? and Lemma 1.2.3, we see that (a, c) is an orbital W of H . By Lemma ??, if (a, c) is an up-bump of f_0 then f_1 moves points near c but not a , and if (a, c) is a down-bump of f_0 then f_1 moves points near a but not c . $\square$

The full converse of the Ubiquity result is false. That is, if $F \cong G \leq H \leq PL_0(\mathbf{I})$, then we cannot reach a conclusion about H as given in Theorem 3.2.

For a counterexample, let x_0 and x_1 be the standard generators of F defined in the introduction. Recall that for all $k > 0$, $x_{k+1} = x_1^{x_0^k}$. Consider the group $H = \langle x_0, f_0, f_1 \rangle$, where $f_0 = x_1^2 x_2^{-1} x_1^{-1}$ and $f_1 = x_1 x_2^2 x_3^{-1} x_2^{-1} x_1^{-1}$.

From Lemma 2.3.4, $\langle f_0, f_1 \rangle \cong F$. From our knowledge about the support of a subgroup (Lemma 1.2.3), we have $(0, 1) = \text{Supp}(x_0) \subseteq \text{Supp}(H) \subseteq (0, 1)$,

which makes $(0, 1)$ the only orbital of H . Let h be any element in H . Since both f_0 and f_1 are the identity near 0, there is an $\varepsilon \in (0, 1)$ and a unique $n \in \mathbb{Z}$ so that $h|_{(0, \varepsilon)} = x_0^n|_{(0, \varepsilon)}$. Similarly, there is a $\delta \in (0, 1)$ and a unique $m \in \mathbb{Z}$ so that $h|_{(\delta, 1)} = x_0^m|_{(\delta, 1)}$. Since the support of x_0 is $(0, 1)$, it must be the case that $n = m$. So then $h|_{(\delta, 1)}$ is the identity if and only if $m = n = 0$ if and only if $h|_{(0, \varepsilon)}$ is the identity. Thus H has no element that moves points near 0 but not near 1, nor vice versa.

References

- [1] Collin Bleak, *a Minimal Non-solvable Group of Homeomorphisms, Groups, Geometry, and Dynamics* **3** (2009), no. 1, 1–37.
- [2] Collin Bleak and Bronlyn Wassink, *Finite Index Subgroups of R. Thompson’s Group F*, arXiv:0711.1014, (2007).
- [3] Matthew G. Brin, *The ubiquity of Thompson’s group F in groups of piecewise linear homeomorphisms of the unit interval*, J. London Math. Soc. (2) **60** (1999), no. 2, 449–460.
- [4] Matthew G. Brin, *Higher dimensional Thompson groups*, Geom. Dedicata **108** (2004), 163–192.
- [5] Matthew G. Brin and Craig C. Squier, *Groups of piecewise linear homeomorphisms of the real line*, Invent. Math. **79** (1985), no. 3, 485–498.
- [6] Matthew G. Brin and Craig C. Squier, *Presentations, conjugacy, roots, and centralizers in groups of piecewise linear homeomorphisms of the real line*, Comm. in Algebra **29** (2001), no. 10, 4557–4596.
- [7] José Burillo, *Quasi-isometrically embedded subgroups of Thompson’s group F*, J. Algebra **212** (1999), 65–78.
- [8] José Burillo, Sean Cleary, and Claas Röver, *Commensurations and Subgroups of Finite Index of Thompson’s Group F*, arXiv:0711.0919 (2007), 1–12.
- [9] J. W. Cannon, W. J. Floyd, and W. R. Parry, *Introductory notes on Richard Thompson’s groups*, Enseign. Math. (2) **42** (1996), no. 3-4, 215–256.